MedStar Health, Inc. POLICY AND PROCEDURE MANUAL

Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

MP.123.MH – Incontinence, Biofeedback

This policy applies to the following lines of business:

✓ MedStar CareFirst PPO

MedStar Health considers **Biofeedback for the treatment of Incontinence** medically necessary for the following indications:

- A. Adult (members 18 years age and older) Biofeedback for Urinary Incontinence and Fecal Incontinence/Constipation when all of the following criteria are met:
- 1. Treatment is billed by a practitioner according to their scope of practice for a diagnosis of urinary incontinence or fecal incontinence/constipation.
- 2. Potential treatable problems should be identified and treatment implemented prior to biofeedback therapy.
- 3. Coverage includes treatment of urinary stress and/or urge incontinence and fecal incontinence/constipation in cognitively intact members.
- 4. Member as failed a documented trial of four weeks of an ordered plan of pelvic muscle exercises (PME), designed to increase periurethral muscle strength.
 - B. Pediatric (members between five years of age and 18 years of age) –Biofeedback for Dysfunctional Elimination Syndrome:

Biofeedback for pediatric members must be prescribed by the urologist and performed by the practitioner according to their scope of practice for the treatment of any of the following conditions associated with bladder dysfunction:

- 1. Nocturnal enuresis
- 2. Staccato voiding
- 3. Bladder-sphincter dyssynergia
- 4. Recurrent urinary tract infections when all of the following criteria are met:
 - a. Physical limitations such as a kidney infection, diabetes, or defects in the urinary system have been ruled out.
 - b. Medications and other conservative measures to address this problem have not reduced or eliminated the issue.
 - c. Dysfunctional elimination (bedwetting) occurs nightly.

Limitations



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

A. Adult (members 18 years age and older) - Biofeedback for Urinary Incontinence and Fecal Incontinence/Constipation:

- 1. Home use of biofeedback therapy is not covered.
- 2. Biofeedback training in a group setting is not covered.
- 3. Biofeedback therapy is limited to up to six treatments per six months per condition.

Appropriate candidates for biofeedback training include:

- Capable of participating in the treatment plan (physically as well as intellectually)
- Motivated to actively participate in the treatment plan, including being responsive to care requirements
- Have a condition that can be appropriately treated with biofeedback.

B. Pediatric (members between five years of age and 18 years of age) – Biofeedback for Dysfunctional Elimination Syndrome

Biofeedback therapy should not be a treatment for any of the following conditions/symptoms and will not be covered:

- 1. This is not a treatment option for children under five years of age.
- 2. Unusual straining during urination or a small or narrow stream of urine or dripping may be a sign of other physical problems.
- 3. Cloudy or pink urine, or bloodstains on underwear or night clothes.

Other limitations include:

- a. Treatment is limited to ten sessions per lifetime.
- b. No coverage provided for the electromyography (EMG) biofeedback device.

Biofeedback therapy for encopresis or constipation in children is considered experimental and investigative.

Background

CMS defines fecal incontinence as the involuntary loss of stool (gas, liquid or solid). Fecal incontinence is caused by a disruption of the normal function of both the lower digestive tract and the nervous system. Fecal incontinence can be caused by several factors:

- Constipation
- Damage to the anal sphincter muscle (e.g., childbirth or hemorrhoid surgery)
- Damage to the nerves of the anal sphincter muscles of the rectum (e.g., childbirth, straining to pass stool, stroke, physical disability due to injury, diabetes or multiple sclerosis)
- Loss of the storage capacity in the rectum



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

- Diarrhea
- Pelvic floor dysfunction

CMS defines urinary incontinence as the involuntary leakage of urine. Male and females have different risk factors in developing urinary incontinence. The risk of urinary incontinence increases with age in both men and women, but women are more likely to develop urinary incontinence due to anatomical differences in the pelvic region and due to changes caused by pregnancy and childbirth. There are several types of urinary incontinence:

- Stress incontinence
- Urge incontinence
- Overflow incontinence
- Mixed incontinence

Biofeedback training of the pelvic muscles uses an electronic or mechanical device to relay visual and/or auditory evidence of pelvic floor muscle activity in order to improve awareness and control of pelvic floor muscle contractions. Biofeedback may include measurement of muscle contraction through surface EMG, vaginal or rectal sensors, and/or rectal manometry devices. Such services when performed for biofeedback monitoring are part of the biofeedback service.

Codes:

Coues.		
CPT Codes / HCPCS Codes / ICD-10 Codes		
Code	Description	
CPT Codes (for Adult (members 18 years age and older) Biofeedback for Urinary Incontinence and Fecal Incontinence/ Constipation):		
90911	Biofeedback training; perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry	
90901	Biofeedback Training by any modality	
CPT codes not covered:		
90875 & 90876	If psychophysiological therapy is performed incorporating biofeedback training, consult these CPT codes	
ICD-10 codes covered if selection criteria are met:		
G83.4	Cauda equine syndrome	
K59.00-K59.09	Constipation	
K59.4	Anal spasm	



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

N31.2	Flaccid neuropathic bladder, not elsewhere classified
N31.9	Neuromuscular dysfunction of bladder
N36.42	Intrinsic sphincter deficiency (ISD)
N36.43	Post-void dribbling
N36.44	Muscular disorders of urethra
G83.4	Cauda equine syndrome
N36.8	Other specified disorders of urethra
N39.3	Stress incontinence (female) (male)
N39.41	Urge incontinence
N39.42	Incontinence without sensory awareness
N39.43	Post-void dribbling
N39.44	Nocturnal enuresis
N39.45	Continuous leakage
N39.46	Mixed incontinence
N39.490	Overflow incontinence
R15.0-R15.9	Fecal incontinence
R33.0	Drug induced retention of urine
R33.8-R33.9	Retention of urine
R35.0	Frequency of micturition
R39.14	Feeling of incomplete bladder emptying
R39.15	Urgency of urination
CPT Codes (for Pediatric (members between five years of age and 18 years of age) Biofeedback for Dysfunctional Elimination Syndrome):	
90911	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry
ICD-10 codes co	vered if selection criteria are met:
N13.70-N13.739	Vesicoureteral reflux
N32.81	Overactive bladder
N36.41-N36.44	Sphincter disorders/ deficiency
N39.0	Urinary tract infection, site not specified
1439.0	ormary tract infoaton, one not opcomed



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

References

- Bassotti G, Chistolini F., Sietchiping-Nzepa F, et al. Biofeedback for pelvic floor dysfunction in constipation. BMJ 2004 Feb; 328(7436): 393-396 http://www.bmj.com/content/328/7436/393?view=long&pmid=14962877

- 5. Desantis DJ, Leonard MP, Preston MA, et al. Effective of biofeedback for dysfunctional elimination syndrome in pediatrics: a systemic review. J Pediatr Urology. 2011 Jun; 7(3): 342-348. http://www.ncbi.nlm.nih.gov/pubmed/21527216
- 6. Hayes Technology Brief. Biofeedback for the Treatment of Dysfunction Voiding in Children. Annual Review March 10, 2011.
- 7. Palmer LS, Biofeedback in the management of urinary continence in children. Curr Urology Rep. 2010 Mar; 11(2): 122-127. http://www.ncbi.nlm.nih.gov/pubmed/20425100
- 8. Porena M., Constantini E., Rociola W, et al. Biofeedback successfully cures detrusor-sphincter dyssynergia in pediatric patients. J.Urol. 2000, June: 162(6); 1927-1931. http://www.ncbi.nlm.nih.gov/pubmed/10799231
- Society of Urological Nurses and Associates. Continence Coalition Statement on Ethical Use of Biofeedback and Electrical Stimulation for the Treatment of Incontinence. Continence Coalition: June 2001. Approved by SUNA Board of Directors. October 2001. Reaffirmed June 2006. http://suna.inurse.com/sites/default/files/download/85-417.pdf



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

- 10. U. S. Department of Health and Human Services (HHS). National Institute for Diabetes and Digestive and Kidney Diseases. Constipation. NIH Publication No. 13-2754. Accessed July 2017. Available at: http://digestive.niddk.nih.gov/ddiseases/pubs/constipation/
- 11.U.S. Department of Health & Human Services (HHS). Agency for Healthcare Research and Quality (AHRQ). National Quality Measures Clearinghouse (NQMC). Management of urinary incontinence in older adults: the percentage of Medicare members 65 years of age and older who reported having a urine leakage problem in the past six months and who discussed their urinary leakage problem with their current practitioner. NQMC #9131. Date of Endorsement: Jan. 3, 2013. Last updated: April 23, 2014. http://www.qualitymeasures.ahrq.gov/content.aspx?id=48665
- 12. Vande WJ, Rittig S, Bauer S, et al. Practical consensus guidelines for the management of enuresis. Eur J Pediatr. 2012 Jun; 171(6): 971-983. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3357467/pdf/431_2012_Article_16 87.pdf
- 13. Weatherall M. Biofeedback or pelvic floor muscle exercises for female genuine stress incontinence. a meta-analysis of trials identified in a systematic review. BJU Int 1999; 83(9): 1015-1016. http://www.ncbi.nlm.nih.gov/pubmed/10368247
- 14. Yagci S., Kibar Y, Akay O, et al. The effect of biofeedback in treatment on voiding and urodynamic parameters in children with voiding dysfunction. J Urol. 2005 November; 174(5):1994 -1998. http://www.ncbi.nlm.nih.gov/pubmed/16217376

Disclaimer:

MedStar Health medical payment and prior authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. The policies constitute only the reimbursement and coverage guidelines of MedStar Health and its affiliated managed care entities. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies.

MedStar Health reserves the right to review and update the medical payment and prior authorization guidelines in its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

These policies are the proprietary information of Evolent Health. Any sale, copying, or dissemination of said policies is prohibited.



Policy Number: MP.123.MH Last Review Date: 08/09/2018 Effective Date: 10/01/2018

